

1-RATIONAL SINGULARITIES AND QUOTIENTS BY REDUCTIVE GROUPS

DANIEL GREB

ABSTRACT. We prove that good quotients of algebraic varieties with 1-rational singularities also have 1-rational singularities. This refines a result of Boutot on rational singularities of good quotients.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

Generalising the Hochster-Roberts theorem [HR74], Jean-François Boutot proved that the class of varieties with rational singularities is stable under taking good quotients by reductive groups, see [Bou87].

In this short note we study varieties with *1-rational singularities*. This is the natural class of singular varieties to which projectivity results for Kähler Moishezon manifolds generalise, cf. [Nam02]. Our main result is:

Theorem 3.1. *Let G be a complex reductive Lie group and let X be an algebraic G -variety such that the good quotient $\pi : X \rightarrow X//G$ exists. If X has 1-rational singularities, then $X//G$ has 1-rational singularities.*

In our proof, which forms a part of the author's thesis [Gre08a], we follow [Bou87] and we check that in Boutot's arguments it is possible to separate the different cohomology degrees.

Theorem 3.1 has been used in [Gre08b] to prove projectivity of compact momentum map quotients of algebraic varieties.

Acknowledgements. The author wants to thank Miles Reid for kindly answering his questions via e-mail. The author gratefully acknowledges the financial support of the Mathematical Sciences Research Institute, Berkeley, by means of a postdoctoral fellowship during the 2009 "Algebraic Geometry" program.

2. PRELIMINARIES

2.1. Singularities and resolution of singularities. We work over the field $\mathbb{C}$ of complex numbers. If X is an algebraic variety we denote by $\mathrm{tdim}_x X$ the dimension of the Zariski tangent space at $x \in X$. I.e., if $\mathfrak{m}_x$ denotes the maximal ideal in $\mathcal{O}_{X,x}$, then $\mathrm{tdim}_x X := \dim_{\mathbb{C}} \mathfrak{m}_x / \mathfrak{m}_x^2$. A point x in an algebraic variety X is called *singular* if $\mathrm{tdim}_x X > \dim X$. We denote the singular set of an algebraic variety X by X_{sing} . A variety X is called *smooth* if $X_{\mathrm{sing}} = \emptyset$.

Date: 22nd January 2009.

Mathematical Subject Classification: 14L30, 14L24, 14B05.

Keywords: group actions on algebraic varieties, good quotient, singularities.

Definition 2.1. Let X be an algebraic variety. A *resolution* of X is a proper birational surjective morphism $f : Y \rightarrow X$ from a smooth algebraic variety Y to X .

If X is an algebraic variety, by a theorem of Hironaka [Hir64] there exists a resolution $f : Y \rightarrow X$ by a projective morphism f such that the restriction $f : f^{-1}(X \setminus X_{\text{sing}}) \rightarrow X \setminus X_{\text{sing}}$ is an isomorphism. See also [BM91], [EH02], and [Kol07] for later improvements and simplifications of Hironaka's proof.

2.2. 1-rational singularities. In this section we introduce the class of singularities studied in this note.

Definition 2.2. An algebraic variety X is said to have *1-rational singularities*, if the following two conditions are fulfilled:

- (1) X is normal,
- (2) for every resolution $f : \tilde{X} \rightarrow X$ of X we have $R^1 f_* \mathcal{O}_{\tilde{X}} = 0$.

Proposition 2.3. Let X be a normal algebraic variety. If there exists one resolution $f_0 : X_0 \rightarrow X$ of X such that $R^1(f_0)_* \mathcal{O}_{X_0} = 0$, then X has 1-rational singularities.

Proof. Let $f_1 : X_1 \rightarrow X$ be a second resolution of X . By [Hir64], there exists a smooth algebraic variety Z and resolutions $g_0 : Z \rightarrow X_0$ and $g_1 : Z \rightarrow X_1$ such that the following diagram commutes

$$\begin{array}{ccc} & Z & \\ g_0 \swarrow & & \searrow g_1 \\ X_0 & & X_1 \\ f_0 \searrow & & \swarrow f_1 \\ & X & \end{array} .$$

For $j = 0, 1$ there exists a spectral sequence (see [Wei94]) with lower terms

$$0 \rightarrow R^1(f_j)_*((g_j)_* \mathcal{O}_Z) \rightarrow R^1(f_j \circ g_j)_* \mathcal{O}_Z \rightarrow (f_j)_*(R^1(g_j)_* \mathcal{O}_Z) \rightarrow \dots$$

Since g_0 and g_1 are resolutions of smooth algebraic varieties, we have $R^1(g_j)_* \mathcal{O}_Z = 0$ and $(g_j)_* \mathcal{O}_Z = \mathcal{O}_{X_j}$ for $j = 0, 1$ (see [Hir64] and [Uen75]). It follows that

$$0 = R^1(f_0)_* \mathcal{O}_{X_0} \cong R^1(f_0 \circ g_0)_* \mathcal{O}_Z \cong R^1(f_1 \circ g_1)_* \mathcal{O}_Z \cong R^1(f_1)_* \mathcal{O}_{X_1}. \quad \square$$

Remark 2.4. The proof shows that, if $f_1 : X_1 \rightarrow X$ and $f_2 : X_2 \rightarrow X$ are two resolutions of X , there is an isomorphism $R^1(f_1)_* \mathcal{O}_{X_1} \cong R^1(f_2)_* \mathcal{O}_{X_2}$. From this, it follows that having 1-rational singularities is a local property. Furthermore, if X is an algebraic G -variety for an algebraic group G , and $f : \tilde{X} \rightarrow X$ is a resolution of X , then the support of $R^1 f_* \mathcal{O}_{\tilde{X}}$ is a G -invariant subvariety of X .

2.2.1. Rational singularities. In this section we shortly discuss the relation of the notion "1-rational singularity" to the more commonly used notion of "rational singularity".

Definition 2.5. An algebraic variety X is said to have *rational singularities*, if the following two conditions are fulfilled:

- (1) X is normal,

(2) for every resolution $f : \tilde{X} \rightarrow X$ of X we have $R^j f_* \mathcal{O}_{\tilde{X}} = 0$ for all $j = 1, \dots, \dim X$.

Remark 2.6. Again, the vanishing of the higher direct image sheaves is independent of the chosen resolution.

Due to a result of Malgrange [Mal57, p. 236] asserting that $R^{\dim X} f_* \mathcal{O}_{\tilde{X}} = 0$ for every resolution $f : \tilde{X} \rightarrow X$ of an irreducible variety X , an algebraic surface has 1-rational singularities if and only if it has rational singularities. These notions differ in higher dimensions as is illustrated by the following example.

Example 2.7. Let Z be a smooth quartic hypersurface in $\mathbb{P}_3$ and let X be the affine cone over Z in $\mathbb{C}^4$. The variety X has an isolated singularity at the origin. Let L be the total space of the line bundle $\mathcal{O}_Z(-1)$, i.e., the restriction of the dual of the hyperplane bundle of $\mathbb{P}_3$ to Z . Then, blowing down the zero section $\mathcal{Z}_L \subset L$ and setting $\tilde{X} := L$, we obtain a map $f : \tilde{X} \rightarrow X$ which is a resolution of singularities, an isomorphism outside of the origin $0 \in X$ with $f^{-1}(0) = \mathcal{Z}_L \cong Z$. We claim that $0 \in X$ is a 1-rational singularity which is not rational.

To see that the origin is a normal point of X it suffices to note that it is obtained as the blow-down of the maximal compact subvariety $\mathcal{Z}_L$ of the smooth variety L .

To compute $(R^j f_* \mathcal{O}_{\tilde{X}})_0$, we use the Leray spectral sequence

$$\dots \rightarrow H^j(X, \mathcal{O}_X) \rightarrow H^j(\tilde{X}, \mathcal{O}_{\tilde{X}}) \rightarrow H^0(X, R^j f_* \mathcal{O}_{\tilde{X}}) \rightarrow H^{j+1}(X, \mathcal{O}_X) \rightarrow \dots$$

and the fact that X is affine to show that $H^j(\tilde{X}, \mathcal{O}_{\tilde{X}}) \cong H^0(X, R^j f_* \mathcal{O}_{\tilde{X}}) = (R^j f_* \mathcal{O}_{\tilde{X}})_0$. Expanding cohomology classes into Taylor series along fibres of L , we get that $H^j(\tilde{X}, \mathcal{O}_{\tilde{X}}) \cong \bigoplus_{k \geq 0} H^j(Z, \mathcal{O}_Z(k))$. Hence, we have

$$(1) \quad (R^j f_* \mathcal{O}_{\tilde{X}})_0 \cong \bigoplus_{k \geq 0} H^j(Z, \mathcal{O}_Z(k)) \quad \text{for all } j \geq 1.$$

It follows from (1) and [Har77, Chap III, Ex 5.5] that $(R^1 f_* \mathcal{O}_{\tilde{X}})_0 = 0$, and hence that $0 \in X$ is a 1-rational singularity. Since the canonical bundle $\mathcal{K}_Z$ of Z is trivial, it follows from Serre duality that $H^2(Z, \mathcal{O}_Z(k)) \cong H^0(Z, \mathcal{O}_Z(-k))$. As a consequence, we get

$$(2) \quad H^2(Z, \mathcal{O}_Z(k)) = \begin{cases} \mathbb{C} & \text{for } k = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Together with (1) this implies that $(R^2 f_* \mathcal{O}_{\tilde{X}})_0 = \mathbb{C}$. Consequently, the singular point $0 \in X$ is not a rational singularity.

2.3. Good quotients.

Definition 2.8. Let G be a complex reductive Lie group acting algebraically on an algebraic variety X . An algebraic variety Y together with a morphism $\pi : X \rightarrow Y$ is called *good quotient* of X by the action of G , if

- (1) π is G -invariant, surjective, and affine,
- (2) $(\pi_* \mathcal{O}_X)^G = \mathcal{O}_Y$.

Example 2.9. Let X be an affine G -variety. Then the algebra of invariants $\mathbb{C}[X]^G$ is finitely generated, and its inclusion into $\mathbb{C}[X]$ induces a regular map π from X to $Y := \text{Spec}(\mathbb{C}[X]^G)$ which fulfills (1) and (2) above, i.e., π is a good quotient.

3. SINGULARITIES OF GOOD QUOTIENTS: PROOF OF THE MAIN RESULT

Let G be a complex reductive Lie group and let X be an algebraic G -variety such that the good quotient $X//G$ exists. We study the singularities of $X//G$ relative to the singularities of X .

More precisely, we prove the following theorem, which is the main result of this note.

Theorem 3.1. *Let G be a complex reductive Lie group and let X be an algebraic G -variety such that the good quotient $\pi : X \rightarrow X//G$ exists. If X has 1-rational singularities, then $X//G$ has 1-rational singularities.*

Before proving the theorem we explain two technical lemmata. The first one discusses the relation between cohomology modules of X and of $X//G$.

Lemma 3.2. *Let X be an algebraic G -variety with good quotient $\pi : X \rightarrow X//G$. Then the natural map $\pi^* : H^1(X//G, \mathcal{O}_{X//G}) \rightarrow H^1(X, \mathcal{O}_X)$ is injective.*

Proof. Let $\mathcal{U} = \{U_i\}_{i \in I}$ be an affine open covering of $X//G$. We can compute the cohomology module $H^1(X//G, \mathcal{O}_{X//G})$ via Čech cohomology with respect to the covering $\mathcal{U}$. Since π is an affine map, $\pi^{-1}(\mathcal{U}) := \{\pi^{-1}(U_i)\}_{i \in I}$ is an affine open covering of X and we can compute the cohomology module $H^1(X, \mathcal{O}_X)$ via Čech cohomology with respect to the covering $\pi^{-1}(\mathcal{U})$.

Let $\eta = (\eta_{ij}) \in C^1(\mathcal{U}, \mathcal{O}_{X//G})$ be a Čech cocycle such that the pullback of the associated cohomology class $[\eta] \in H^1(X//G, \mathcal{O}_{X//G})$ fulfills $\pi^*([\eta]) = 0 \in H^1(X, \mathcal{O}_X)$. Then, there exists a cocycle $\nu = (\nu_i) \in C^0(\pi^{-1}(\mathcal{U}), \mathcal{O}_X)$ such that

$$\pi^*(\eta_{ij}) = \nu_i|_{\pi^{-1}(U_{ij})} - \nu_j|_{\pi^{-1}(U_{ij})} \in \mathcal{O}_X(\pi^{-1}(U_{ij})).$$

Averaging $\nu_i \in \mathcal{O}_X(\pi^{-1}(U_i))$ over a maximal compact subgroup K of G we obtain invariant functions $\tilde{\nu}_i \in \mathcal{O}_X(\pi^{-1}(U_i))^G$. Since $\pi^*(\eta_{ij}) \in \mathcal{O}_X(\pi^{-1}(U_{ij}))^G$, the cocycle $\tilde{\nu} = (\tilde{\nu}_i) \in C^0(\pi^{-1}(\mathcal{U}), \mathcal{O}_X)$ fulfills

$$\pi^*(\eta_{ij}) = \tilde{\nu}_i|_{\pi^{-1}(U_{ij})} - \tilde{\nu}_j|_{\pi^{-1}(U_{ij})} \in \mathcal{O}_X(\pi^{-1}(U_{ij})).$$

For all i , there exist a uniquely determined function $\hat{\nu}_i \in \mathcal{O}_{X//G}(U_i)$ with $\pi^*(\hat{\nu}_i) = \tilde{\nu}_i$. Consequently, we have

$$\eta_{ij} = \hat{\nu}_i|_{U_{ij}} - \hat{\nu}_j|_{U_{ij}} \in \mathcal{O}_{X//G}(U_{ij}).$$

Therefore, $[\eta] = 0 \in H^1(X//G, \mathcal{O}_{X//G})$ and π^* is injective, as claimed. $\square$

The second lemma will be used to obtain information about the singularities of an algebraic variety X from information about the singularities of a general hyperplane section H of X and vice versa.

Lemma 3.3. *Let X be a normal affine variety with $\dim X \geq 2$ and let $f : \tilde{X} \rightarrow X$ be a resolution of singularities. Let $\mathcal{L} \subset \mathcal{O}_X(X)$ be a finite-dimensional subspace, such that the associated linear system is base-point free, and such that the image of X under the associated map $\varphi_{\mathcal{L}} : X \rightarrow \mathbb{P}_N$ fulfills $\dim(\varphi_{\mathcal{L}}(X)) \geq 2$. If $h \in \mathcal{L}$ is a general element, then the following holds for the corresponding hyperplane section $H \subset X$:*

- (1) *The preimage $\tilde{H} := f^{-1}(H)$ is smooth and $f|_{\tilde{H}} : \tilde{H} \rightarrow H$ is a resolution of H .*
- (2) *We have $R^j f_* \mathcal{O}_{\tilde{H}} \cong R^j f_* \mathcal{O}_{\tilde{X}} \otimes \mathcal{O}_H$ for $j = 0, 1$.*

Proof. 1) This follows from Bertini's theorem (see [Har77, Chap III, Cor 10.9]).

2) In the exact sequence

$$(3) \quad 0 \rightarrow \mathcal{O}_{\tilde{X}}(-\tilde{H}) \xrightarrow{m} \mathcal{O}_{\tilde{X}} \rightarrow \mathcal{O}_{\tilde{H}} \rightarrow 0,$$

the map m is given by multiplication with the equation $h \in \mathcal{O}_X(X)$ defining H and $\tilde{H}$. Pushing forward the short exact sequence (3) by f_* yields the long exact sequence

$$(4) \quad \begin{aligned} 0 \rightarrow f_*\mathcal{O}_{\tilde{X}}(-\tilde{H}) &\xrightarrow{m_0} f_*\mathcal{O}_{\tilde{X}} \rightarrow f_*\mathcal{O}_{\tilde{H}} \rightarrow \\ &\rightarrow R^1f_*\mathcal{O}_{\tilde{X}}(-\tilde{H}) \xrightarrow{m_1} R^1f_*\mathcal{O}_{\tilde{X}} \rightarrow R^1f_*\mathcal{O}_{\tilde{H}} \rightarrow \\ &\rightarrow R^2f_*\mathcal{O}_{\tilde{X}}(-\tilde{H}) \xrightarrow{m_2} R^2f_*\mathcal{O}_{\tilde{X}} \rightarrow R^2f_*\mathcal{O}_{\tilde{H}} \rightarrow \\ &\rightarrow \dots \end{aligned}$$

Since X is an affine variety, the exact sequence above is completely determined by the following sequence of finite $\mathcal{O}_X(X)$ -modules:

$$(5) \quad \begin{aligned} 0 \rightarrow \Gamma(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\tilde{H})) &\xrightarrow{m_0} \Gamma(\tilde{X}, \mathcal{O}_{\tilde{X}}) \rightarrow \Gamma(\tilde{X}, \mathcal{O}_{\tilde{H}}) \rightarrow \\ &\rightarrow H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\tilde{H})) \xrightarrow{m_1} H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}) \rightarrow H^1(\tilde{X}, \mathcal{O}_{\tilde{H}}) \rightarrow \\ &\rightarrow H^2(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\tilde{H})) \xrightarrow{m_2} H^2(\tilde{X}, \mathcal{O}_{\tilde{X}}) \rightarrow H^2(\tilde{X}, \mathcal{O}_{\tilde{H}}) \rightarrow \\ &\rightarrow \dots \end{aligned}$$

The maps $m_j, j = 0, 1, 2$ are given by multiplication with the element $h \in \mathcal{O}_X(X)$. We claim that we can choose $h \in \mathcal{L}$ in such a way that m_1 and m_2 are injective. Indeed, if R is a commutative Noetherian ring with unity, M is a finite R -module, and $Z_R(M)$ denotes the set of zerodivisors for M in R , we have

$$Z_R(M) = \bigcup_{P \in \text{Ass } M} P,$$

where $\text{Ass } M$ is the finite set of assassins (or associated primes) of M . Hence, the set of zerodivisors for $H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\tilde{H}))$ and $H^2(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\tilde{H}))$ is a union of finitely many prime ideals P of $\mathcal{O}_X(X)$. Since the linear system associated to $\mathcal{L}$ is base-point free, the general element h of $\mathcal{L}$ lies in $\mathcal{L} \setminus \bigcup P$. For such a general $h \in \mathcal{L} \setminus \bigcup P$, the maps m_1 and m_2 in the sequences (5) and (4) are injective.

Since $\mathcal{O}_{\tilde{X}}(-\tilde{H}) \cong f^*(\mathcal{O}_X(-H))$, the projection formula for locally free sheaves (see [Har77, Chap III, Ex 8.3]) yields

$$R^j f_* \mathcal{O}_{\tilde{X}}(-\tilde{H}) \cong R^j f_* \mathcal{O}_{\tilde{X}} \otimes \mathcal{O}_X(-H).$$

Furthermore, for $j = 0, 1$ the image of m_j coincides with the image $\mathcal{B}_j$ of the natural map $R^j f_* \mathcal{O}_{\tilde{X}} \otimes \mathcal{O}_X(-H) \rightarrow R^j f_* \mathcal{O}_{\tilde{X}}$. Since m_1 and m_2 are injective by the choice of H , it follows that

$$R^j f_* \mathcal{O}_{\tilde{H}} \cong R^j f_* \mathcal{O}_{\tilde{X}} / \mathcal{B}_j \quad \text{for } j = 0, 1.$$

Tensoring with $R^j f_* \mathcal{O}_{\tilde{X}}$ is right-exact, and hence the exact sequence

$$0 \rightarrow \mathcal{O}_X(-H) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_H \rightarrow 0$$

yields $R^j f_* \mathcal{O}_{\tilde{H}} \cong R^j f_* \mathcal{O}_{\tilde{X}} \otimes \mathcal{O}_H$, as claimed. $\square$

Proof of Theorem 3.1. Since the claim is local and π is an affine map, we may assume that $X//G$ and X are affine.

First, we prove that normality of X implies normality of $X//G$. We have to show that $\mathbb{C}[X//G] \cong \mathbb{C}[X]^G$ is a normal ring. So let $\alpha \in \text{Quot}(\mathbb{C}[X]^G) \subset \mathbb{C}(X)^G$ be an element of the quotient field of $\mathbb{C}[X]^G$ and assume that α fulfills a monic equation

$$\alpha^n + c_1\alpha^{n-1} + \cdots + c_n = 0$$

with coefficients $c_j \in \mathbb{C}[X]^G \subset \mathbb{C}[X]$. Since $\mathbb{C}[X]$ is normal by assumption, it follows that $\alpha \in \mathbb{C}(X)^G \cap \mathbb{C}[X] = \mathbb{C}[X]^G$. Hence, $\mathbb{C}[X]^G$ is normal. As a consequence, we can assume in the following that X is G -irreducible.

We prove the claim by induction on $\dim X//G$. For $\dim X//G = 0$ there is nothing to show. For $\dim X//G = 1$ we notice that $X//G$ is smooth. So, let $\dim X//G \geq 2$. Let $\pi : X \rightarrow X//G$ denote the quotient map and let $p_X : \tilde{X} \rightarrow X$ be a resolution of X . First, we prove that a general hyperplane section $H \subset X//G$ has 1-rational singularities. If H is a general hyperplane section in $X//G$, Lemma 3.3 applied to $\pi^{-1}(H)$ yields that $p_X|_{\tilde{H}} : \tilde{H} \rightarrow \pi^{-1}(H)$ is a resolution, where $\tilde{H} = p_X^{-1}(\pi^{-1}(H))$, and that

$$R^j(p_X)_* \mathcal{O}_{\tilde{X}} \otimes_{\mathcal{O}_X} \mathcal{O}_{\pi^{-1}(H)} = R^j(p_X)_* \mathcal{O}_{\tilde{H}} \quad \text{for } j = 0, 1.$$

Since $f_* \mathcal{O}_{\tilde{X}} = \mathcal{O}_X$ by Zariski's main theorem, it follows from the case $j = 0$ that $\pi^{-1}(H)$ is normal. Alternatively, one could invoke Seidenberg's Theorem (see e.g. [BS95, Thm. 1.7.1]). Together with the case $j = 1$, this implies that $\pi^{-1}(H)$ has 1-rational singularities. By induction, it follows that $H = \pi^{-1}(H)//G$ has 1-rational singularities.

Let $p : Z \rightarrow X//G$ be a resolution of $X//G$. As we have seen above, a general hyperplane section H of $X//G$ has 1-rational singularities and the restriction of p to $\hat{H} := p^{-1}(H)$, $p|_{\hat{H}} : \hat{H} \rightarrow H$ is a resolution of H . It follows that $\mathcal{O}_H \otimes R^1 p_* \mathcal{O}_Z = R^1 p_* \mathcal{O}_{\hat{H}} = 0$. Consequently, the support of $R^1 p_* \mathcal{O}_Z$ does not intersect H and hence, $\text{supp}(R^1 p_* \mathcal{O}_Z)$ consists of isolated points. Since the claim is local, we can assume in the following that $R^1 p_* \mathcal{O}_Z$ is supported at a single point $x_0 \in X//G$.

The group G acts on the fibre product $Z \times_{X//G} X$ such that the map $p_X : Z \times_{X//G} X \rightarrow X$ is equivariant. One of the G -irreducible components $\tilde{X}$ of $Z \times_{X//G} X$ is birational to X , and, by passing to a resolution of $\tilde{X}$ if necessary, we can assume that $p_X : \tilde{X} \rightarrow X$ is a resolution of X . We obtain the following commutative diagram

$$\begin{array}{ccc} X & \xleftarrow{p_X} & \tilde{X} \\ \pi \downarrow & & \downarrow p_Z \\ X//G & \xleftarrow{p} & Z. \end{array}$$

Since $R^1 p_* \mathcal{O}_Z$ is supported only at x_0 , we have $(R^1 p_* \mathcal{O}_Z)_{x_0} = H^0(X//G, R^1 p_* \mathcal{O}_Z)$. Recall that $X//G$ is affine, hence, the Leray spectral sequence

$$0 \rightarrow H^1(X//G, \mathcal{O}_{X//G}) \rightarrow H^1(Z, \mathcal{O}_Z) \rightarrow H^0(X//G, R^1 p_* \mathcal{O}_Z) \rightarrow H^2(X//G, \mathcal{O}_{X//G}) \rightarrow \cdots$$

implies that it suffices to show that $H^1(Z, \mathcal{O}_Z) = 0$.

Since X is affine and has 1-rational singularities, it follows from the Leray spectral sequence

$$0 \rightarrow H^1(X, \mathcal{O}_X) \rightarrow H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}) \rightarrow H^0(X, R^1 f_* \mathcal{O}_{\tilde{X}}) \rightarrow \dots$$

that $H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}) \cong H^1(X, \mathcal{O}_X) = 0$. Consequently, it suffices to show that there exists an injective map $H^1(Z, \mathcal{O}_Z) \hookrightarrow H^1(\tilde{X}, \mathcal{O}_{\tilde{X}})$.

We introduce the following notation: $U = (X//G) \setminus \{x_0\}$, $U' = \pi^{-1}(U) \subset X$, $\tilde{U} = p_X^{-1}(U') \subset \tilde{X}$, $V = p^{-1}(U) \subset Z$. We obtain the following commutative diagram of canonical maps

$$(6) \quad \begin{array}{ccccc} H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}) & \xrightarrow{h_{\tilde{X}, \tilde{U}}} & H^1(\tilde{U}, \mathcal{O}_{\tilde{U}}) & \xleftarrow{h_{U', \tilde{U}}} & H^1(U', \mathcal{O}_{U'}) \\ \uparrow h_{Z, \tilde{X}} & & \uparrow & & \uparrow h_{U, U'} \\ H^1(Z, \mathcal{O}_Z) & \xrightarrow{h_{Z, V}} & H^1(V, \mathcal{O}_V) & \xleftarrow{h_{U, V}} & H^1(U, \mathcal{O}_U). \end{array}$$

Let $Y := p^{-1}(x_0) \subset Z$. If $H_Y^1(Z, \mathcal{O}_Z)$ denotes the local cohomology groups with support in Y , there exists an exact sequence

$$(7) \quad \dots \rightarrow H_Y^1(Z, \mathcal{O}_Z) \rightarrow H^1(Z, \mathcal{O}_Z) \xrightarrow{h_{Z, V}} H^1(V, \mathcal{O}_V) \rightarrow \dots,$$

see [Har77, Chap III, Ex 2.3]. We use the following vanishing result which is originally due to Hartshorne and Ogus [HO74].

Proposition 3.4. *Let W be a normal affine algebraic variety and $p : Z \rightarrow W$ a resolution. Let $w \in W$ and $Y = p^{-1}(w)$. Then $H_Y^i(Z, \mathcal{O}_Z) = 0$ for all $i < \dim W$.*

Sketch of proof. Set $n := \dim W$. As a first step, we compactify p . There exist projective completions $\bar{Z}$ and $\bar{W}$ of Z and W , respectively, and a resolution $\bar{p} : \bar{Z} \rightarrow \bar{W}$ such that $\bar{p}^{-1}(W) = Z$ and $\bar{p}|_Z = p$. Since Z is an open neighbourhood of Y in $\bar{Z}$, the Excision Theorem of Local Cohomology (see [Har77, Chap III, Ex 2.3]) implies that $H_Y^j(\bar{Z}, \mathcal{O}_{\bar{Z}}) = H_Y^j(Z, \mathcal{O}_Z)$. Let $\mathcal{K}_{\bar{Z}}$ be the locally free sheaf associated to the canonical bundle of $\bar{Z}$. The Formal Duality Theorem (see [Har70]) implies that the dual $H_Y^j(\bar{Z}, \mathcal{O}_{\bar{Z}})^*$ of $H_Y^j(\bar{Z}, \mathcal{O}_{\bar{Z}})$ is isomorphic to $(R^{n-j} \bar{p}_* \mathcal{K}_{\bar{Z}})_{\bar{w}}^\wedge$, where $^\wedge$ denotes completion with respect to the maximal ideal of $\mathcal{O}_{\bar{W}, w} = \mathcal{O}_{W, w}$. In summary, we have obtained an isomorphism

$$H_Y^j(Z, \mathcal{O}_Z)^* \cong (R^{n-j} \bar{p}_* \mathcal{K}_{\bar{Z}})_{\bar{w}}^\wedge \quad \text{for all } j = 0, \dots, n.$$

By Grauert-Riemenschneider vanishing (see e.g. [Laz04, Chap 4.3.B]), the term on the right hand side equals zero for $n - j \geq 1$. This proves the claim. $\square$

Since $\dim X//G \geq 2$, Proposition 3.4 yields $H_Y^1(Z, \mathcal{O}_Z) = 0$. As a consequence of (7), the map $h_{Z, V}$ is injective.

The restriction of p to $V = p^{-1}(U)$ is a resolution of U . Since the support of $R^1 p_* \mathcal{O}_Z$ is concentrated at x_0 , the variety U has 1-rational singularities, and the Leray spectral sequence

$$0 \rightarrow H^1(U, \mathcal{O}_U) \xrightarrow{h_{U, V}} H^1(V, \mathcal{O}_V) \rightarrow H^0(U, R^1 p_* \mathcal{O}_U) \rightarrow \dots$$

yields that $h_{U, V}$ is bijective. Similar arguments show that $h_{U', \tilde{U}}$ is bijective. Furthermore, Lemma 3.2 implies that $h_{U, U'}$ is injective.

By the considerations above, the map

$$h_{Z,\tilde{U}} := h_{U',\tilde{U}} \circ h_{U,U'} \circ h_{U,V}^{-1} \circ h_{Z,V}$$

is injective. Diagram (6) implies $h_{Z,\tilde{U}} = h_{\tilde{X},\tilde{U}} \circ h_{Z,\tilde{X}}$, and therefore $h_{Z,\tilde{X}}$ is injective. Consequently, we have $H^1(Z, \mathcal{O}_Z) = 0$. This concludes the proof of Theorem 3.1. $\square$

REFERENCES

- [BM91] Edward Bierstone and Pierre D. Milman, *A simple constructive proof of canonical resolution of singularities*, Effective methods in algebraic geometry (Castiglione, 1990), Progr. Math., vol. 94, Birkhäuser, Boston, MA, 1991, pp. 11–30.
- [Bou87] Jean-François Boutot, *Singularités rationnelles et quotients par les groupes réductifs*, Invent. Math. **88** (1987), no. 1, 65–68.
- [BS95] Mauro C. Beltrametti and Andrew J. Sommese, *The adjunction theory of complex projective varieties*, de Gruyter Expositions in Mathematics, vol. 16, Walter de Gruyter & Co., Berlin, 1995.
- [EH02] Santiago Encinas and Herwig Hauser, *Strong resolution of singularities in characteristic zero*, Comment. Math. Helv. **77** (2002), no. 4, 821–845.
- [Gre08a] Daniel Greb, *Projectivity of analytic Hilbert quotients*, Dissertation, Ruhr-Universität Bochum, 2008, available at: <http://www-brs.ub.ruhr-uni-bochum.de/netahtml/HSS/Diss/GrebDaniel/diss.pdf>.
- [Gre08b] ———, *Projectivity of analytic Hilbert quotients*, <http://arxiv.org/abs/0805.3973>, 2008.
- [Har70] Robin Hartshorne, *Ample subvarieties of algebraic varieties*, Lecture Notes in Mathematics, vol. 156, Springer-Verlag, Berlin, 1970.
- [Har77] ———, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer-Verlag, New York, 1977.
- [Hir64] Heisuke Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II*, Ann. of Math. (2) **79** (1964), 109–203; *ibid.* (2) **79** (1964), 205–326.
- [HO74] Robin Hartshorne and Arthur Ogus, *On the factoriality of local rings of small embedding codimension*, Comm. Algebra **1** (1974), 415–437.
- [HR74] Melvin Hochster and Joel L. Roberts, *Rings of invariants of reductive groups acting on regular rings are Cohen-Macaulay*, Advances in Math. **13** (1974), 115–175.
- [Kol07] János Kollár, *Lectures on resolution of singularities*, Annals of Mathematics Studies, vol. 166, Princeton University Press, Princeton, NJ, 2007.
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, vol. 48, Springer-Verlag, Berlin, 2004.
- [Mal57] Bernard Malgrange, *Faisceaux sur des variétés analytiques réelles*, Bull. Soc. Math. France **85** (1957), 231–237.
- [Nam02] Yoshinori Namikawa, *Projectivity criterion of Moishezon spaces and density of projective symplectic varieties*, Internat. J. Math. **13** (2002), no. 2, 125–135.
- [Uen75] Kenji Ueno, *Classification theory of algebraic varieties and compact complex spaces*, Lecture Notes in Mathematics, vol. 439, Springer-Verlag, Berlin, 1975.
- [Wei94] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.

MATHEMATISCHES INSTITUT, ABTEILUNG FÜR REINE MATHEMATIK, ALBERT-LUDWIGS-UNIVERSITÄT, ECKERSTR. 1, 79104 FREIBURG IM BREISGAU, GERMANY

E-mail address: daniel.greb@math.uni-freiburg.de

URL: <http://home.mathematik.uni-freiburg.de/dgreb>